

Foliar Fungicides for Control of Corn Diseases

The following information has been developed on fungicides for management of certain foliar diseases of corn for use in Ohio.

Fungicides for corn disease control based on appropriate application timing.

Product	Fungicide(s)	Rate/A (fl. oz.)	Gray leaf spot	Anthrachnose leaf blight	Northern corn leaf blight	Eye spot	Common rust	PHI
Strobilurin	Headline 2.09 EC	Pyraclostrobin 23.6%	6.0 - 12.0	x	x	x	x	7 days
	Quadris 2.08 SC	Azoxystrobin 22.9%	6.0 - 15.5	x	x	x	x	7 days
Triazole	Folicur 3.6 F	Tebuconazole 38.7%	4.0 - 6.0	x		x	x	36 days
	PropiMax 3.6 EC	Propiconazole 41.8%	4.0	x		x	x	30 days
	Tilt 3.6 EC	Propiconazole 41.8%	2.0 - 4.0	x		x	x	30 days
Premix	Quilt 200 SC	Azoxystrobin 7.0% Propiconazole 11.7%	7.0 - 14.0	x		x	x	30 days
	Stratego 250 EC	Propiconazole 11.4% Trifloxystrobin 11.4%	10.0 -12.0	x	x	x	x	30 days

This information is provided only as a guide. It is the responsibility of the pesticide applicator by law to read and follow all current label directions. No endorsement is intended for products listed, nor is criticism meant for products not listed.