

Wheat Seed Treatment Fungicides for Ohio

The following information has been developed on fungicides for management of certain seedling diseases of wheat in Ohio.

Fungicides for wheat seed treatment

Product	Fungicide(s)	Rate/100 lb seed	Loose smut	Common Bunt	Stagonospora	Fusarium (scab)	Pythium
Allegiance FL	Metalaxyl 28.35%	0.375 - 0.75 fl oz	N	N	N	N	E
Apron XL	Mefenoxam 33.3%	0.32 - 0.64 fl oz	N	N	N	N	E
Charter	Triticonazole 2.4%	3.1 fl oz	E	E		G-E	P
Dividend Extreme	Difenoconazole 7.7% Mefenoxam 1.93%	2 - 4 fl oz	E	E	E	G	E
Incentive RTA	Difenoconazole 3.21% Mefenoxam 0.27%	2.5 – 10 fl oz	E	E		G	E
Rancona Pinnacle	Ipconazole 0.434% Matalaxyl 0.579%	5.0 – 8.33 fl oz					
Maxim 4FS	Fludioxonil 40.3%	0.08 – 0.16 fl oz	N	N	N	G	N
Raxil XT	Metalaxyl 20% Tebuconazole 15%	0.16 fl oz	E	E	E	G	E
Raxil-Thiram	Thiram 20% Tebuconazole 0.6%	3.5 – 4.6 fl oz	E	E	E	G	F
Vitavax	Various formulations	Refer to product label	G	G	F	G	F

Efficacy based on labeled rates of active ingredient for each material.

Efficacy rating scale: E=excellent, G=good, F=fair, P=poor, N=no activity, ND=no data

This information is provided only as a guide. It is the responsibility of the pesticide applicator by law to read and follow all current label directions.

No endorsement is intended for products listed, nor is criticism meant for products not listed.