



Bayer CropScience LP  
 P.O. Box 12014, 2 T.W. Alexander Drive  
 Research Triangle Park, North Carolina 27709  
 1-866-99BAYER (1-866-992-2937)  
<http://www.bayercropscienceus.com>

**Folicur® 3.6 F Foliar Fungicide**

**For Use on Soybean to control Rust  
 (*Phakopsora pachyrhizi*)**

**SECTION 18 SPECIFIC EXEMPTION  
 File Symbol: 04-OH-10**

**FOR DISTRIBUTION AND USE ONLY IN THE STATE OF OHIO  
 UNDER SECTION 18 SPECIFIC EXEMPTION**

This labeling must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the registered product label for Folicur® 3.6 F Foliar Fungicide (EPA Registration Number 264-752).

Any adverse effects resulting from the use of Folicur® 3.6 F Foliar Fungicide under this specific exemption must be immediately reported to the State of Ohio Department of Agriculture.

**EFFECTIVE PERIOD:**

This labeling becomes effective 08/27/04 and expires on 03/01/07.

**DIRECTIONS FOR USE**

A maximum of 4.3 million acres of soybeans may be treated under this exemption.

**RECOMMENDED APPLICATIONS**

CROP	DISEASES	RATE OF FOLICUR® 3.6 F
Soybean	Rust ( <i>Phakopsora pachyrhizi</i> )	3 to 4 fl oz per acre

**Use Directions:** Apply Folicur® 3.6 F as a broadcast foliar spray at first appearance of rust pustules on any leaves within the plant canopy. Repeat applications on a 10 to 14-day spray interval if environmental conditions are favorable for continued disease development. Applications may be made up to the R6 (full seed) soybean growth stage. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. For optimum benefit, the lowest label recommended rate of a spray surfactant must be tank-mixed with Folicur® 3.6 F. Folicur® 3.6 F should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.

**Restrictions:** Applications may not be made within 21 days of harvest. Do not apply more than 2 applications per season. Do not apply more than 8 fl. oz/A per use season. Do not enter treated fields for 12 hours after application. Do not feed soybean forage or hay to livestock.

**ENVIRONMENTAL HAZARDS**

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

For **MEDICAL** And **TRANSPORTATION** Emergencies **ONLY** Call 24 Hours A Day 1-800-334-7577

For **PRODUCT USE** Information Call 1-866-99BAYER (1-866-992-2937)

or

Visit our worldwide web site at [www.bayercropscienceus.com](http://www.bayercropscienceus.com)

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).