

# SUPPLEMENTAL LABELING

## DUPONT™ UPPERCUT™ FUNGICIDE

**DuPont Crop  
Protection**

### DUPONT™ UPPERCUT™ FUNGICIDE

EPA Reg. No. 264-752-352

#### FOR USE ON SOYBEAN TO CONTROL RUST

*(Phakopsora pachyrhizi)*

##### SECTION 18 SPECIFIC EXEMPTION

**File Symbol: 04-OH-10**

**FOR DISTRIBUTION AND USE ONLY IN THE  
STATE OF OHIO**

**UNDER SECTION 18 SPECIFIC EXEMPTION**

This labeling must be in possession of the user at the time of pesticide application.

Any adverse effects resulting from the use of UPPERCUT™ fungicide under this specific exemption must be immediately reported to the State of Ohio Department of Agriculture.

##### EFFECTIVE PERIOD:

This label becomes effective 06/30/05 and expires on 11/10/07.

A maximum of 3 total applications using approved Section 18 products collectively will be made under this soybean rust Section 18.

A maximum of 4.3 million acres of soybeans may be treated under this exemption.

##### ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment washwater or rinsate.

##### DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

##### RECOMMENDED APPLICATIONS

Crop	Diseases	Rate of UPPERCUT™
Soybean	Rust <i>(Phakopsora pachyrhizi)</i>	3 to 4 fl oz per acre

**Use Directions:** Apply UPPERCUT™ as a broadcast foliar spray prior to or at first appearance of rust pustules on any leaves within the plant canopy. Repeat applications on a 10- to 21-day spray interval if environmental conditions are favorable for continued disease development. Use of the higher rates and shorter spray intervals are recommended when disease pressure is severe. Use of adjuvants may enhance the performance of UPPERCUT™. If utilized, apply the lowest labeled rate of the adjuvant to enhance disease control. UPPERCUT™ should be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 5 gallons per acre by aircraft spray equipment.

**Restrictions:** Applications may not be made within 30 days of harvest. Do not apply more than 8 fl. oz/A per use season. Do not enter treated fields for 12 hours after application. Do not feed soybean forage or hay to livestock. Do not make more than 2 applications of tebuconazole-containing products to soybeans per season. Do not apply more than 0.225 pounds of the active ingredient tebuconazole per acre per year. A maximum of three total applications using approved section 18 products collectively are allowed under this soybean rust section 18.

© 2005 E. I. du Pont de Nemours and Company, Crop Protection, Wilmington, Delaware 19898

**IMPORTANT  
BEFORE USING THIS PRODUCT FUNGICIDE, READ  
AND CAREFULLY NOTE THE PRECAUTIONARY  
STATEMENTS AND OTHER PROCEDURAL INFORMA-  
TION APPEARING ON THE EPA REGISTERED LABEL  
OR ON OTHER SUPPLEMENTAL LABELS.**

This bulletin contains new or supplemental instructions for use of these products in combination which does not appear on the package label. Follow the instructions carefully.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

R-505 070105 063005